

MYCOTAXON

<http://dx.doi.org/10.5248/126.231>

Volume 126, pp. 231–233

October–December 2013

***Flavophlebia sphaerospora*,
a new corticoid species from India**

MANINDER KAUR, AVNEET P. SINGH * & G.S. DHINGRA

*Department of Botany, Punjabi University, Patiala 147002, India**CORRESPONDENCE TO: avneetbot@gmail.comABSTRACT — *Flavophlebia sphaerospora* sp. nov. is described from Himachal Pradesh, India.KEY WORDS — *Basidiomycota*, Northwestern Himalaya, Narkanda

During the fungal forays conducted during August 2012 in the Hattu Peak area, Narkanda, district Shimla of Himachal Pradesh, India, Maninder and Avneet collected a fungus on decaying wood of *Cedrus deodara* in a mixed forest. After comparison of macroscopic and microscopic characters (Parmasto 1968, Hjortstam & Larsson 1977, Bernicchia & Gorjón 2010, Mycobank 2013) we concluded that the material represented the monotypic genus *Flavophlebia* but differed from *F. sulfureoisabellina* (Litsch.) K.H. Larss. & Hjortstam.

***Flavophlebia sphaerospora* Man. Kaur, Avneet P. Singh & Dhingra, sp. nov. PLATE 1**

MYCOBANK 804821

Differs from *Flavophlebia sulfureoisabellina* by its denser subiculum and its spherical to subspherical basidiospores.

TYPE: India, Himachal Pradesh: Shimla, Narkanda, on way to Hattu Peak, on the bark of decaying wood of *Cedrus deodara* (Roxb. ex D. Don) G. Don, 19 August 2012, Maninder Kaur & Avneet 5166 (PUN, **holotype**).

ETYMOLOGY: The epithet refers to the shape of the basidiospores.

Basidiocarp resupinate, adnate, effused, ceraceous, ≤ 180 μm thick in section; hymenial surface smooth, grayish green to grayish yellow to grayish orange; margins thinning, paler concolorous, to indeterminate. Hyphal system monomitic; generative hyphae branched, septate, clamped; basal hyphae ≤ 3.3 μm wide, more or less parallel to the substrate, thick-walled, forming a dense subiculum; subhymenial hyphae ≤ 2.7 μm wide, vertical, densely united.





PLATE 1. *Flavophlebia sphaerospora* (holotype). 1. Basidiocarp showing hymenial surface. 2. Vertical section through basidiocarp showing hyphae, basidia, cystidia and basidiospores.

Cystidia 45–60 × 8–8.5 µm, cylindrical, flexuous, thin-walled, with basal clamp. Basidia 28–36 × 8.5–10 µm, clavate, somewhat stalked, 4-sterigmate, with basal clamp, filled with oily contents; sterigmata ≤6.0 µm long. Basidiospores 6.6–8.0 µm in diameter, spherical to subspherical with subapical apiculus reminding of *Radulomyces confluens* (Fr.) M.P. Christ., with numerous oil-droplets, thin- to somewhat thick-walled, inamyloid, somewhat cyanophilous.

REMARKS— Both *Flavophlebia sphaerospora* and *F. sulfureoisabellina* occur on the bark of coniferous trees and have a similar greenish-yellowish colour, but *F. sulfureoisabellina* differs in having a comparatively loose subiculum, ellipsoid to subglobose basidiospores, and slightly narrower basidia.

Acknowledgements

The authors thank Head (Department of Botany, Punjabi University, Patiala) for providing research facilities, Dr. Nils Hallenberg (Professor Emeritus in Botany, Goteborg, Sweden) for peer review and expert comments, and Prof. B.M. Sharma (Department of Plant Pathology, COA, CSKHPAU, Palampur, H.P., India) for peer review.

Literature cited

- Bernicchia A. & Gorjón SP. 2010. *Corticiaceae* s.l. Fungi Europaei 12. Edizioni Candusso. Alassio. Italia. 1008 p.
- Hjortstam K. & Larsson, KH. 1977. Notes on *Corticiaceae* (*Basidiomycetes*). Mycotaxon 5(2): 475–480.
- Mycobank. 2013. Fungal databases. Nomenclature and species banks. [Accessed: 11/07/2013] <http://www.mycobank.org/>
- Parmasto E. 1968. Conspectus Systematis Corticiacearum. Tartu. 262 p.